

## **MIRALAX BOWEL PREP**

### **SUPPLIES TO PURCHASE**

Purchase the following medications at least 2 days prior to your procedure. If you have kidney issues or cannot afford the prep, call our office.

- Two bottles of Miralax 238g (powder)
- Dulcolax 5mg tablets (bisacodyl)
- Two 32 ounce bottles of Gatorade or Powerade (no red) or Crystal Light

### **5 DAYS BEFORE PROCEDURE**

Stop foods that contain skins, hulls, seeds and nuts (peanuts, popcorn, grapes, beans, peels of apples/potatoes, strawberries, tomatoes, bread with seeds, rice, etc.). No salads.

### **1 DAY BEFORE PROCEDURE**

- **No** solid foods
- **No** dairy products
- **No** red, orange, or purple liquids (Jell-O, sports drinks, juices or popsicles). Other colors are okay.
- Start clear liquid diet. Examples include: juice (white grape, white cranberry, lemonade), clear broth, soda (7-UP, Coke, etc.), black coffee and tea, popsicles, plain Jell-O (no fruit chunks), sports drinks, and Ensure Clear.

**8AM** – Take 4 Dulcolax tablets with 8 ounces of water

**6PM** – Mix one bottle of Miralax powder into 32oz of Gatorade. Drink ENTIRE bottle of Gatorade/Miralax mixture over the next hour (8oz every 15 minutes).

### **DAY OF PROCEDURE**

#### **6 Hours before procedure**

- Mix one bottle of Miralax powder into 32oz of Gatorade.
- Drink ENTIRE bottle of Gatorade/Miralax mixture over the next hour (8oz every 15 minutes).
- **DO NOT DRINK ANYTHING ONCE FINISHED WITH GATORADE/MIRALAX**

It is normal to feel abdominal cramping and bloating and have multiple loose stools during the prep period. Nausea and vomiting may also occur. Complete all of the prep if possible. A successful prep is when your bowel movements are a **clear, yellow liquid**.

\*\*If you have any questions, call 812-882-2703 and ask for a nurse.